

**Chronic Joint Symptoms - (Calculated Variable)*
Kentucky BRFSS 2001**

Geographic Groups	Total Respond.#	Yes			No		
		N	%	C.I.(95%)	N	%	C.I.(95%)
TOTAL	7493	2607	30.2	(28.8-31.7)	4886	69.8	(68.3-71.2)
ADDs							
Barren River	529	189	32.0	(27.8-36.6)	340	68.0	(63.4-72.2)
Big Sandy	432	190	40.6	(35.4-46.0)	242	59.4	(54.0-64.6)
Bluegrass	525	163	29.3	(24.9-34.2)	362	70.7	(65.8-75.1)
Buffalo Trace	420	134	29.7	(22.6-38.0)	286	70.3	(62.0-77.4)
Cumberland Valley	464	181	37.2	(32.3-42.4)	283	62.8	(57.6-67.7)
FIVCO	560	225	37.6	(33.0-42.3)	335	62.4	(57.7-67.0)
Gateway	578	196	30.5	(25.9-35.5)	382	69.5	(64.5-74.1)
Green River	413	140	29.7	(25.1-34.8)	273	70.3	(65.2-74.9)
Kentucky River	482	213	38.2	(33.3-43.3)	269	61.8	(56.7-66.7)
KIPDA	560	160	26.0	(22.2-30.2)	400	74.0	(69.8-77.8)
Lake Cumberland	487	170	30.4	(26.0-35.2)	317	69.6	(64.8-74.0)
Lincoln Trail	497	154	28.2	(24.1-32.8)	343	71.8	(67.2-75.9)
Northern Kentucky	553	162	28.0	(24.1-32.4)	391	72.0	(67.6-75.9)
Pennyrile	441	147	30.3	(25.9-35.2)	294	69.7	(64.8-74.1)
Purchase	552	183	30.5	(26.4-35.0)	369	69.5	(65.0-73.6)

*The data are calculated from 'yes' responses to two questions:1)During the past twelve months have you had pain or stiffness in or around a joint?2)Were these symptoms present on most days for at least one month?

Response is generalized to the state population

#Use caution in interpreting data for total respondents less than 50. N = Cell Size, NH = Non-Hispanic, %

= Percentage, C.I.(95%) = Confidence Interval (at 95 percent probability level).

Percentages are weighted to population characteristics.

Unless stated differently above, Missing, 'Don't know', and Refused responses are excluded.